

Berry Crop Protection Update

These lists include new products and product changes for the 2026 season. Check out the [Crop Protection Hub](#) or the [label \(PMRA label search\)](#) for information on rate, preharvest interval, restricted entry interval, efficacy, and comments.

Click the link in the ‘**Crop**’ column to see the product in the Crop Protection Hub.

Label expansions & new products in 2025-2026

Product, a.i., Group, PCP	Crop	Pests	Comments
Insecticides & Miticides			
Magister <i>Fenazaquin</i> Group 21 (34544)	Raspberry	Two-spotted spider mite	Rate: 1.75-2.34 L/ha. Apply in at least 500 L/ha. Do not apply when bees are active.
	Blueberry	Blueberry bud mite	Rate: 1.75 L/ha. Apply in at least 500 L/ha. This pest has not been detected in Ontario. Do not apply when bees are active.
Spear T <i>GS-omega/kappa-Hxtx-Hv1a</i> Group 32 (34679)	Raspberry Blueberry	Spotted wing drosophila (suppression only)	Rate is 9.4-28 L/ha. Use with a spreader/sticker or non-ionic surfactant at 0.125% where permitted.
Fungicides			
Orondis Gold <i>Metalaxyl-M and S-isomer + oxathiapiprolin</i> Group 4 + 49 (33508)	Strawberry	Phytophthora crown rot (suppression only) ; Red Stele	Apply in a minimum of 2,000 L water/ha.
	Raspberry Blueberry	Phytophthora root rot	Apply in a minimum of 2,000 L water/ha.

Product, a.i., Group, PCP	Crop	Pests	Comments
EcoSwing <i>Swinglea glutinosa</i> extract Group BM1 (35206)	Blueberry	Botrytis grey mould (Partial suppression only) ; Mummy berry	0 day PHI
	Raspberry Gooseberry Strawberry Currants	Botrytis grey mould (Partial suppression only)	
LifeGard WG <i>Bacillus mycooides</i> Group P6 (32526)	Blueberry	Botrytis grey mould (Suppression only) ; Anthracnose; Mummy berry	This product works by activating the plant's defense mechanism and is most effective when it is applied before infection occurs.
	Strawberry	Anthracnose; Botrytis grey mould (suppression only) ; powdery mildew (suppression only) ; common leaf spot (suppression only)	
	Raspberry Blackberry Currants Gooseberries	Botrytis grey mould (suppression only)	
Buran Garlic powder Group NC (30601)	Blackberry Raspberry	Fire blight	Rate: 1.4% v/v. For increased coverage, use with a registered non-ionic surfactant.
Cyclone Plus Citric and lactic acid Group NC (34762)	Blueberry	Anthracnose fruit rot	Use a spray volume of 350 – 1000 L/ha.
Salibro <i>Fluazaindolizine</i> Group NC (34182)	Strawberry	Root-knot nematode (suppression only)	Application methods include pre-plant incorporated, chemigation (pre-plant, at-plant, or post-plant), and soil drench (pre-plant, at-plant, or post-plant).

Product, a.i., Group, PCP	Crop	Pests	Comments
Tivano Plus Citric and lactic acid Group NC (34763)	Strawberry	Angular leaf spot, powdery mildew (suppression only)	Do not use silicone-based surfactant if angular leaf spot disease is active.
Gatten <i>Flutianil</i> Group U13 (34297)	Strawberry	Powdery mildew	Apply in 470-1870 L water/ha. The registrant recommends the addition of a registered surfactant to improve coverage. Contact Cohort Wholesale for more information.
Herbicides			
Lontrel XC <i>Clopyralid</i> Group 4 (32795)	Raspberry Blackberry	Suppression or control of Labelled Weeds	Apply Lontrel as a banded application. 30 day PHI.
Eragon <i>Saflufenacil</i> Group 14 (31469)	Blueberry	Post-emerge spot or band application.	Add Merge Adjuvant at a rate of 1 L/ha. Use as a directed spray. Do not allow spray drift to contact dormant canes as severe crop injury will result.

Recent Re-evaluation Decisions for Berry Growers

Registration cancellations		
Product (PCP #)	Crop (Pest)	Decision
Fenbuconazole: Indar (27294)	Blueberry (mummy berry)	Fenbuconazole discontinued. Last date of use was December 31, 2025.

Additionally, production and sale of **Poast Ultra is discontinued** for all use patterns. **Betamix registration has been cancelled**- the last date of use was December 31, 2025. Botector and Apollo are no longer available and have been removed from the Berry sections of the Hub.